

## Yield and Nutritive Value of Anti-Taurus Mountain Rangeland Shrubs in Turkey

<sup>1</sup>Kagan Kokten, <sup>2</sup>Oktay GURSOY, <sup>3</sup>Tuncay TUKEL and <sup>3</sup>Rustu HATİPOĞLU

<sup>1</sup>Department of Field Crops, Faculty of Agriculture, Bingöl University, 12000 Bingöl, Turkey

<sup>2</sup>Department of Animal Science, <sup>3</sup>Department of Field Crops,  
Faculty of Agriculture, Cukurova University, 01330 Adana, Turkey

**Abstract:** Shrubs provide the bulk of feed for goats in the Mediterranean agropastoral farming systems, yet quantitative data on productivity and nutritive value of highland shrubs species that is needed to develop rangeland management strategies are limited. Edible Fodder (EF) yield and concentrations of CP, ADF, NDF and IVOMD in the EF of shrubs were evaluated at Saksak and Candir in Yayladagi County of Antakya on the Anti-Taurus Mountains in Turkey. Significant variations in EF yield (8-4003 kg DM ha<sup>-1</sup>), CP (5.9-23.1%), ADF (19.9-38.4%), NDF (30.9-54.2%) and IVOMD (22.7-56.7%) were recorded among the shrubs at Saksak. Similarly, variations in EF yield (202-1523 kg DM ha<sup>-1</sup>), CP (5.9-13.2%), ADF (22.8-37.7%), NDF (38.3-53.6%) and IVOMD (29.3-51.4%) were significant among the shrubs at Candir. Based on digestible organic matter yield, the shrubs could be ranked as: *Daphne sericea*>*Calycotome infesta*>*Styrax officinalis*>*Cistus salviaefolius*>*Rhamnus alaternus*>*Laurus nobilis*>*Juniperus oxycedrus*>*Quercus coccifera*>*Pistacia terebentis* in Saksak and *Daphne sericea*>*Styrax officinalis*>*Cistus salviaefolius*>*Juniperus oxycedrus*>*Quercus coccifera*>*Pistacia terebentis* in Candir. *D. sericea*, *S. officinalis*, *C. salviaefolius* and *C. infesta* showed greater potential for development of agroforestry technologies to increase rangeland productivity in the Anti-Taurus Mountains region and similar highland environments in other parts of the world. Further research on cheaper methods of establishment, defoliation options to optimize and sustain edible fodder and wood production, fodder quality in terms of outputs of livestock products may be needed.

**Key words:** Chemical composition, edible fodder, eastern mediterranean shrubs, goats, highland, *in vitro* digestibility, Turkey

### INTRODUCTION

The grazing lands of the Taurus Mountains are subject to heavy, uncontrolled grazing pressure and the forage production capacities of these lands are gradually decreasing. The reason for this retrogressive trend includes a complex mixture of several interrelated problems of socio-economic, judiciary, political and technical inadequacies (Tukel and Hatipoglu, 1998).

Located on the Anti-Taurus Mountains in Turkey, Yayladagi County of Antakya has limited land, water and capital resources. As a result, livelihoods of the majority of small-scale farmers in the region depend on rangeland based livestock production using crosses of local (Hair) and Damascus (Shami) goats for milk, meat and manure (Haenlein, 1998; Boyazoglu and Morand-Fehr, 2001; GURSOY, 2005; Mavrogenis, 2005) and cropping of tobacco and wheat with manure as the main nutrient source. Recent reduction in state purchased tobacco to only 200 kg per household has further reduced household

incomes. Hence, most farmers perceive increase in production of livestock products as the fastest way of compensating for the lost incomes, if seasonal feed deficits that constrain increased livestock production could be reduced by improving the degraded rangelands which provide the bulk of feed (Le Houerou, 1986; Papanastasis, 2000).

Goats are reported to be preferably better browsers of shrub vegetation than grazers of herbaceous vegetation (Nastis, 1996; Rogosic *et al.*, 2006).

Although, there have been several notable studies on Taurus Mountain rangelands (Tukel, 1984; Ozer and Tukel, 1988; Genekan and Celen, 1994; Heil *et al.*, 1999; Cinar, 2001), they primarily describe the general range vegetation and forage producing capacities but not the feeding values under different ecological conditions in Turkey. Most of these studies are unavailable to the interested foreign readers. There have also been extensive works on flora of Anti-Taurus Mountains (Yildiz, 1982; Cakan, 1997). Recently, just a few research works

appeared dealing with the nutritive values of some mediterranean shrub and tree species (Kamalak *et al.*, 2004; Karabulut *et al.*, 2006). However, quantitative data is limited on the productivity and nutritive value of Mediterranean highland shrub species, especially those preferred by goats (Le Houerou, 1986; Papachristou, 2000; Aharon *et al.*, 2007; Ainalis *et al.*, 2006). Also, intra-species and inter-species variations in fodder yield and quality have not been widely reported. Such information is needed to develop rangeland improvement plans and agroforestry technologies and to prioritize shrub research (Le Houerou 1986, 1994). These reports on a study to quantify edible fodder yield and nutritional value of shrubs in the Anti-Taurus Mountains region in Turkey.

## MATERIALS AND METHODS

**Study sites:** The experiment was conducted from 2002-2003 at two locations, namely Saksak (1750 m asl, 35°58' 34.55"N and 36°05'37.58"E) and Candir (1235 m asl, 35°56'13.30"N and 36°00'49.22"E) in Yayladagi County of Antakya on the Anti-Taurus Mountains. Yayladagi County is representative of the Taurus Mountains region in the Mediterranean Coast of Turkey. Annual rainfall average 750 mm, humidity 75%, with minimum and maximum temperatures of 6 and 33°C, respectively over a 30 years period. Monthly maximal temperatures were lower but minimal temperatures were higher during the experimental period. More precipitation received in February, August and December of the experimental period than the long term monthly precipitation in the study area. However, there was no significant variation for the relative humidity.

**Estimating edible fodder yield:** Data were collected from three sites at Saksak and two sites in Candir. At each site, three plots (replications) of size 5×10 m were randomly selected and marked. Shrub species in each replicate were identified. Shrub biomass was measured using the reference unit method (Andrew *et al.*, 1979) to estimate the biomass of each shrub individually and then the total biomass of the plantation. Biomass was estimated at three monthly intervals on: 6 April, 8 July and 13 October 2002 and 15 January 2003 by 4 persons. Thereafter, leaves and twigs <10 mm in diameter (Edible Fodder, EF) were separated from the sample branch and weighed fresh. The samples were oven-dried at 70°C for 48 h to determine Dry Matter (DM) content. Total EF yield was estimated from the sum of the four harvests.

**Chemical composition and *in vitro* digestibility analysis:** The oven-dried EF samples for each shrub species were bulked by replicate at each site and ground through

a 1 mm screen to determine total Nitrogen (N) by the Kjeldahl method (AOAC, 1990). Crude Protein (CP) was calculated as N×6.25. Neutral Detergent Fibre (NDF) and Acid Detergent Fibre (ADF) were analyzed according to Van Soest *et al.* (1991). *In vitro* Organic Matter Digestibility (IVOMD) was determined by a modified two stage procedure (Moore and Mott, 1974).

**Statistical analysis:** Data were analyzed separately for Saksak and Candir using the General Linear Models (GLM) procedures (SAS Institute, 1990). Edible fodder yield data were analyzed as a split-plot experiment with site as main-plots and shrubs as sub-plots, whilst the fodder quality data were analyzed as a randomized block experiment using sites as replicates. Univariate correlation analyses were used to establish relationships between concentrations of CP, ADF, NDF and IVOMD in the edible fodder. When significance is not given in the text, it implies to minimum ( $p < 0.05$ ).

## RESULTS AND DISCUSSION

**Fodder yield:** Edible fodder varied significantly among the shrubs at both locations, with yield at Saksak (Fig. 1a) being relatively higher than Candir (Fig. 1b). At Saksak, yield varied significantly from 8 kg DM ha<sup>-1</sup> for *C. salviaefolius* to 4003 kg DM ha<sup>-1</sup> for *Q. coccifera*; whilst at Candir, yield varied from 202 kg DM ha<sup>-1</sup> for *D. sericea* to 1523 kg DM ha<sup>-1</sup> for *Q. coccifera*.

**Crude protein concentration:** Table 1 presents the nutritive value of the EF. The CP concentration ranged from 5.9% in *J. oxycedrus* at both locations to 23.1% in *C. infesta* at Saksak and 13.2% in *P. terebentis* at Candir.

**Cell wall concentration:** At Saksak, concentrations of ADF ranged significantly from 19.9% for *C. infesta* to 38.4% for *Q. coccifera*, whilst NDF ranged from 30.9% for *C. infesta* to 54.2% for *Q. coccifera* (Table 1). At Candir, the ADF concentration varied significantly among the shrubs but the NDF did not differ. *P. terebentis* had the lowest ADF (22.8%) and *Q. coccifera* the highest (37.7%).

***In vitro* organic matter digestibility:** At the Saksak, the IVOMD concentration ranged from as low as 22.7% for *P. terebentis* to 56.7% for *D. sericea*; whilst at the Candir it ranged from 29.3% for *P. terebentis* to 51.4% for *D. sericea*.

**Correlations between determinants of fodder quality:** Positive and significant correlations were recorded between concentrations of CP and IVOMD (Saksak,




Fig. 1: Edible fodder yield of rangeland shrub species at Saksak (a) and Candir (b) in Yayladagi County of Antakya, Turkey

Table 1: Chemical composition and *in vitro* organic matter digestibility of edible fodder of rangeland shrub species at Saksak and Candir in Yayladagi County of Antakya, Turkey

Locations	Shrubs	<sup>1</sup> CP	<sup>1</sup> ADF	<sup>1</sup> NDF	<sup>1</sup> IVOMD	
		----- (%) -----				
Saksak	<i>Calycotome infesta</i>	23.10	19.90	30.90	54.50	
	<i>Cistus salviaefolius</i>	10.20	24.00	43.80	41.80	
	<i>Daphne sericea</i>	11.00	26.70	36.80	56.70	
	<i>Juniperus oxycedrus</i>	5.90	36.30	41.30	32.60	
	<i>Laurus nobilis</i>	10.70	33.00	48.40	35.40	
	<i>Pistacia terebentis</i>	11.60	26.00	36.10	22.70	
	<i>Quercus coccifera</i>	7.90	38.40	54.20	31.60	
	<i>Rhamnus alaternus</i>	9.60	29.30	41.60	40.70	
	<i>Styrax officinalis</i>	10.60	23.90	40.20	46.80	
	Mean	11.18	28.61	41.48	40.31	
	LSD (p<0.05)	2.72	4.46	7.67	9.16	
	Minimum	5.90	19.90	30.90	22.70	
	Maximum	23.10	36.30	54.20	56.70	
	Candir	<i>Cistus salviaefolius</i>	10.90	33.60	49.20	41.30
		<i>Daphne sericea</i>	9.60	29.90	40.80	51.40
<i>Juniperus oxycedrus</i>		5.90	36.30	41.70	35.60	
<i>Pistacia terebentis</i>		13.20	22.80	38.30	29.30	
<i>Quercus coccifera</i>		8.50	37.70	53.60	29.60	
<i>Styrax officinalis</i>		11.60	27.20	43.10	45.70	
Mean		10.00	31.30	44.50	38.80	
LSD (p<0.05)		2.22	7.56	16.11	5.24	
Minimum		5.90	22.80	38.30	29.30	
Maximum		13.20	37.70	53.60	51.40	

<sup>1</sup>CP: Crude Protein; ADF: Acid Detergent Fibre; NDF: Neutral Detergent Fibre; IVOMD: *In vitro* Organic Matter Digestibility

r = 0.61; Candir, r = 0.46) and concentrations of ADF and NDF (Saksak = 0.51). In contrast, negative and significant correlation were recorded between concentrations of CP and ADF (Saksak, r = -0.75; Candir, r = -0.72), ADF and IVOMD (Saksak, r = -0.62; Candir, r = -0.65) and NDF and IVOMD (Saksak, r = -0.47). In agreement with the results, significant variations in EF yield of shrubs have been

reported (Le Houerou, 1986, 1994; Larbi *et al.*, 2007). The differences in yield may relate to variations in anatomical, physiological and morphological characteristics associated with the acquisition of moisture, lights and nutrients for biomass production (Le Houerou, 1986, 1994; Dong and Zhang, 2001). They could also reflect differences among the shrub in residual buds, leaf area index and storage of carbohydrate reserves after defoliation.

Ranges in CP concentration in the current study are consistent with those reported for several shrubs (Le Houerou, 1994; Papachristou, 2000; Haddi *et al.*, 2003; Ammar and Gonzalez, 2005). The significant differences in CP concentration among the shrubs could be partly due to differences in leaf: twig ratio in the samples used for the chemical analysis, cell wall concentration and thickening (Wilson, 1994). All but *J. oxycedrus* and *Q. coccifera* had CP concentration >8.0%, below which digestibility is depressed and voluntary forage intake drops (Minson, 1990). This suggests that most of the shrubs could effectively provide supplemental N, when grazed with native pastures or when offered to stall-fed ruminants fed basal diets of low N cereal crop residues. The availability of the N, however, will depend on the concentration of secondary compounds such as condensed tannins which influence adhesion, colonization and enzymatic activity of the rumen microbial ecosystem and consequently the degradation of the different dietary fractions (Guimaraes-Beelen *et al.*, 2006).

The significant differences in EF cell-wall concentrations among the shrubs confirm earlier reports (Papachristou, 2000; Haddi *et al.*, 2003; Ventura *et al.*, 2004; Larbi *et al.*, 2007) possibly due to anatomical or morphological differences related to cell wall rigidity (Wilson, 1994) and leaf:twig ratio in the samples used for the chemical analyses. The observed differences in ADF and NDF concentrations among the shrubs could have implications for their use as fodder, because cell wall concentration in EF of shrubs is negatively correlated with palatability, voluntary DM intake and potential DM degradability (Kaitho *et al.*, 1998; Larbi *et al.*, 1998).

Significant differences among the shrubs in the IVOMD concentration confirm earlier reports (Le Houerou, 1986; Papachristou, 2000; Larbi *et al.*, 2007). The observed differences in the CP, ADF and NDF concentrations (Table 1) and secondary compounds such as tannin in the EF (Ventura *et al.*, 2004) may be partly accountable. It could also be due to differences in the configuration of cell-wall polysaccharides and their effect on rumen microbial attachment and colonization of digesta particles (Cheng *et al.*, 1984). Larbi *et al.* (2005, 2007) reported significant positive and negative correlations between the determinants of shrub EF which agree with

the results. The significant correlations between CP and IVOMD and ADF and IVOMD suggest that IVOMD could be predicted from either CP or ADF under situations where facilities for determining IVOMD are not available. Similarly, the significant correlation between CP and ADF indicates that ADF could be predicted from CP.

High milk production in the Damascus x Kil Goats found on Anti-Taurus Mountains (Acuz, 2005) could be attributed to the rich *Q. coccifera* composition of the rangelands. As far as range management concerns, higher yields together with slightly higher CP and IVOMD of *Q. coccifera* and *J. oxycedrus* in the Saksak site indicate that more animal units can be stocked in the Saksak rangeland than the Candir site. It is reported that *Q. coccifera* can meet the nutritional requirements of goats all year round but requires supplementation in winter (Papanastasis and Liacos, 1980).

### CONCLUSION

The research priorities in order to reach sound management practices should be focused on determining correct carrying capacities of the study sites as well as following appropriate grazing periods in the area.

### ACKNOWLEDGEMENTS

The researchers would like extend their gratitude to Cukurova University Research Fund and ICARDA (International Center for Agricultural Research in the Dry Areas, Aleppo, Syria) and Antoine Makzume Ltd. Co. of Iskenderun, Antakya, for co-funding of the project. They are also indebted to Mrs. Monika Zaklouta of ICARDA for the analyses of the samples.

### REFERENCES

AOAC., 1990. Official Methods of Analysis. 15th Edn., Association of Official Analytical Chemists, Washington, DC., USA., pp: 200-210.

Acuz, S., 2005. Characterization of the goat breed in the Yayladagi county of Hatay province and determination of variation in growth and milk production. M.Sc. Thesis, University of Cukurova, Institute of Science, Adana, Turkey.

Aharon, H., Z. Henkin, E.D. Ungar, D. Kababya, H. Baram and A. Perevolotsky, 2007. Foraging behavior of the newly introduced Boer goat breed in a Mediterranean woodland: A research observation. *Small Rumin. Res.*, 69: 144-153.

Ainalis, A.B., C.N. Tsiouvaras and A.S. Nastis, 2006. Effect of summer grazing on forage quality of woody and herbaceous species in a silvopastoral system in Northern Greece. *J. Arid Environ.*, 67: 90-99.

Ammar, H. and J.S. Gonzalez, 2005. Assessment of the digestibility of some Mediterranean shrubs by *in vitro* techniques. *Anim. Feed Sci. Technol.*, 119: 323-331.

Andrew, M.H., I.R. Noble and R.T. Lange, 1979. A non-destructive method for estimating the weight of forage on shrubs. *Aust. Rangeland J.*, 1: 225-231.

Boyazoglu, J. and P. Morand-Fehr, 2001. Mediterranean dairy sheep and goat products and their quality: A critical review. *Small Rumin. Res.*, 40: 1-11.

Cakan, H., 1997. Plant ecology of Musa and Keldag Mountains. Ph.D. Thesis, Cukurova University, Institute of Science, Adana, Turkey.

Cheng, K.J., C.S. Stewart, D. Dinsdale and J.W. Costerlon, 1984. Electron bacteria involved in the digestion of plant cell walls. *Anim. Feed Sci. Technol.*, 10: 93-120.

Cinar, S., 2001. A study on the productivity and botanical composition of rangelands in Hanyeri Village of Tufanbeyli County of Adana. M.Sc. Thesis, University of Çukurova, Institute of Science, Adana, Turkey.

Dong, X. and X. Zhang, 2001. Some observation of the adaptations of sandy shrubs to the arid environment in the Mu Us Sandland: Leaf water relations and anatomic features. *J. Arid Environ.*, 48: 41-48.

Gençkan, M.S. and A.E. Celen, 1994. Some floristic Classification researches on the improvement of the maqui-vegetation pastures of Turkey. *Turk. J. Agric. For.*, 18: 303-312.

Guimaraes-Beelen, B., T.T. Berchielli, R. Beelen and A.N. Medeiros, 2006. Influence of condensed tannins from Brazilian semi-arid legumes on ruminal degradability, microbial colonization and ruminal enzymatic activity in Saanen goats. *Small Rumin. Res.*, 61: 35-44.

Gursoy, O., 2005. Small Ruminant Breeds of Turkey. In: Characterization of Small Ruminant Breeds in West Asia and North Africa, Iniguez, L. (Ed.). International Center for Agricultural Research in the Dry Areas, Aleppo, Syria, pp: 239-416.

Haddi, M.L., S. Filacorda, K. Meniai, F. Rollin and P. Susmel, 2003. *In vitro* fermentation kinetics of some halophyte shrubs sampled at three stages of maturity. *Anim. Feed Sci. Technol.*, 104: 215-225.

Haenlein, G.F.W., 1998. The value of goats and sheep to sustain mountain farmers. *Int. J. Anim. Feed Sci.*, 13: 187-194.

Heil, G.W., T. Tukul and M.S.A. Warger, 1999. Regenerative potential of the land use systems in Goksu, Taurus, Turkey. Report on Project Nr SA. 5-2-05 (CRG. 970624) 1096/97/JARC-501. Sponsored by NATO.

- Kaitho, R.J., I.V. Nsahlai, B.A. Williams, N.N. Umunna, S. Tamminga and J. van Bruchem, 1998. Relationships between preference, rumen degradability, gas production and chemical composition of browses. *Agrofor. Syst.*, 39: 129-144.
- Kamalak, A., O. Canbolat, O. Ozay and S. Akbas, 2004. Nutritive value of oak (*Quercus* sp.) leaves. *Small Rumin. Res.*, 53: 161-165.
- Karabulut, A., O. Canbolat, C.O. Ozkan and A. Kamalak, 2006. Potential nutritive value of some Mediterranean shrub and tree leaves as emergency food for sheep in winter. *Livest. Res. Rural Dev.*, 18: 78-87.
- Larbi, A., A. Khatib, A. Bolus, J. Tiedeman and L. Iniguez, 2007. Forage Potential of Leguminous and Non-leguminous Fodder Shrubs in the Dry Areas of Central and West Asia and North Africa. In: Human and Nature-Working Together for Sustainable Development of Drylands, El-Beltagy, A., M.C. Saxena and T. Wang (Eds.). ICARDA., Aleppo, Syria, pp: 468-477.
- Larbi, A., J.W. Smith, I.O. Kurdi, I.O. Adekunle, A.M. Raji and D.O. Ladipo, 1998. Chemical composition, rumen degradation and gas production characteristics of some multipurpose fodder trees and shrubs during wet and dry seasons in humid tropics. *Anim. Feed Sci. Technol.*, 72: 81-96.
- Larbi, A., N.J. Anyanwu, U.I. Oji, I. Etela, L.D. Gbaraneh and D.O. Ladipo, 2005. Fodder yield and nutritive value of browse species in West African humid tropics: response to age of coppice regrowth. *Agrofor. Syst.*, 65: 197-205.
- Le Houerou, H.N., 1986. Salt tolerant plants of economic value in the Mediterranean basin. *Reclamation Revegetation Res.*, 5: 319-341.
- Le Houerou, H.N., 1994. Forage Halophytes and Salt tolerant Fodder Crops in the Mediterranean Basin. In: Halophytes as a Resource for Livestock and for Rehabilitation of Degraded Lands, Squires, V.R. and A.T. Ayoub (Eds.). Kluwer Academic Publishers, Netherlands, pp: 124-137.
- Mavrogenis, A.P., 2005. Small Ruminant Breeds of Cyprus. In: Characterization of Small Ruminant Breeds in West Asia and North Africa, Iniguez, L. (Ed.). ICARDA., Aleppo, Syria, pp: 417-458.
- Minson, D.J., 1990. Forages in Ruminant Nutrition. Academic Press, New York, pp: 350.
- Moore, J.E. and G.O. Mott, 1974. Recovery of residual organic matter from *in vitro* digestion of forages. *J. Dairy Sci.*, 57: 1258-1280.
- Nastis, A., 1996. Feeding behaviour of goats and utilization of pasture and rangelands. Proceedings of the International Conference on Goats, (ICG'96), International Academic Publishers, Beijing, China, pp: 487-494.
- Ozer, A. and T. Tukul, 1988. A study on the assessment of productivity and botanical composition of grazed and protected rangelands in Kesmeburun Village of Osmaniye. *C.U. Inst. Sci. J. Sci. Eng.*, 2: 21-35.
- Papachristou, T.G., 2000. Nutritive Value of Fodder Shrubs use in Mediterranean Sylvo-pastoral Systems and their use by Grazing Animals. In: Fodder Shrub Development in Arid and Semi-Arid Zones, Gintzburger, G., M. Bounejmate and N. Nefzaoui (Eds.). Vol. 2, ICARDA., Aleppo, Syria, pp: 461-468.
- Papanastasis, V.P. and L.G. Liacos, 1980. Productivity and management of kermes oak brushlands for goats. Proceedings of the International Symposium on Browse in Africa, ILCA., Addis Ababa.
- Papanastasis, V.P., 2000. Shrubland Management and Shrub Plantations in Southern Europe. In: Fodder Shrub Development in Arid and Semi-Arid Zones, Gintzburger, G., M. Bounejmate and N. Nefzaoui (Eds.). Vol. 1, ICARDA., Aleppo, Syria, pp: 54-66.
- Rogosic, J., J.A. Pfister, F.D. Provenza and D. Grbesa, 2006. Sheep and goat preference for and nutritional value of mediterranean maquis shrubs. *Small Ruminant Res.*, 64: 169-179.
- SAS Institute, Inc., 1990. SAS User's Guide. Statistical Analysis System Institute, Cary, NC, USA.
- Tukul, T. and R. Hatipoglu, 1998. Grazing management problems and their consequences in Turkey. Proceedings of the International Workshop on Ecological Basis of Livestock Grazing in Mediterranean Ecosystems, Oct. 23-25, Thessaloniki, Greece, pp: 318-321.
- Tukul, T., 1984. Comparison of grazed and protected mountain steppe rangeland in Ulukisla, Turkey. *J. Range Manage.*, 37: 133-135.
- Van Soest, P.J., J.B. Robertson and B.A. Lewis, 1991. Methods for dietary fiber, neutral detergent fiber and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.*, 74: 3583-3597.
- Ventura, M.R., J.I.R. Castanon, M.C. Pieltain and M.P. Flores, 2004. Nutritive value of forage shrubs: *Bituminaria bituminosa*, *Rumex lunaria*, *Acacia salicina*, *Cassi sturtii* and *Adenocarpus foliosus*. *Small Rumin. Res.*, 52: 13-18.
- Wilson, J.R., 1994. Cell wall characteristics in relation to forage digestion by ruminants. *J. Agric. Sci.*, 122: 173-182.
- Yildiz, B., 1982. A study on the flora of Berit Mountain (Kahramanmaras). Ph.D. Thesis, Hacettepe University Faculty of Education, Department of Botanic, Ankara.